

The following is an extract from:

Nutrient Reference Values for Australia and New Zealand
Including Recommended Dietary Intakes

ENDORSED BY THE NHMRC ON 9 SEPTEMBER 2005

© Commonwealth of Australia 2006

ISBN Print 1864962372
ISBN Online 1864962437

The Nutrient Reference Values (NRVs) was a joint initiative of the Australian National Health and Medical Research Council (NHMRC) and the New Zealand Ministry of Health (MoH). The NHMRC would like to thank the New Zealand MoH for allowing the use of the NRV material in the development of this website.

NHMRC publications contact:

Email: nhmrc.publications@nhmrc.gov.au
Internet: <http://www.nhmrc.gov.au>
Free Call: 1800 020 103 ext 9520

FOLATE

BACKGROUND

Folate is the commonly used group name for folic acid (pteroyl glutamic acid, or PGA) and its derivatives with similar activity. In foods and in the body folates are usually in the reduced form (tetrahydrofolate, or THF) and conjugated with up to seven glutamate residues and one of several types of one-carbon groups. PGA is used in supplements and for food fortification as it is more stable than the other derivatives.

Folate functions as a coenzyme in single-carbon transfers in the metabolism of nucleotides and amino acids. It is essential for the formation of thymidylate (TMP) for DNA synthesis, so that without folate, living cells cannot divide. The need for folate is higher when cell turnover is increased, such as in fetal development. It is also involved in purine synthesis, in the generation of formate and in amino acid interconversions. Homocysteine is methylated by methyl-THF to produce methionine, which is in turn used for the synthesis of *S*-adenosyl-methionine an important methylating agent *in vivo* (Wagner 1996).

Food folates are hydrolysed to monoglutamate forms in the gut to allow their absorption across the intestine. The monoglutamates enter the portal circulation and are metabolised to polyglutamate derivatives in the liver. They are either retained, or released to the blood as reconverted monoglutamates or to bile. The liver contains about 50% of the body stores of folate.

Folate is a substrate and vitamin B₁₂ is a coenzyme for the formation of MTHF that depends on the regeneration of THF, the parent compound in the homocysteine-to-methionine conversion. If either folate or vitamin B₁₂ is deficient, megaloblastic changes occur in bone marrow and other replicating cells from lack of 5,10-MTHF for DNA synthesis.

The bulk of excretion products are folate cleavage products. Intact urinary folate accounts for only a small percentage of dietary folate. Biliary excretion of folate can be as high as 100 µg/day (Herbert & Das 1993, Whitehead 1986), however much of this is reabsorbed.

Folate is difficult to measure in foods because it is present in different forms, so food databases can be inaccurate. However, the main sources of folate in Australia and New Zealand according to the National Nutrition Surveys undertaken in 1995 and 1997, respectively (ABS 1998, MOH 1999), are cereals, cereal products and dishes based on cereals (about 27%) and vegetables and legumes (about 29%). Fruit provides about 8–10%. Orange juice is contributing a greater amount than in the past due to the recent introduction of fortification with folate.

Folate requirements can be affected by bioavailability, nutrient interactions, smoking, certain drugs and genetic variations. Notably, the C667T polymorphism that causes MTHF reductase deficiency is found in 2–16% of white populations (van der Put et al 1995). It is likely that individuals who are homozygous for this polymorphism may have a higher requirement for folate.

Bioavailability of folates in food is about 50–60% whereas that of the folic acid used to fortify foods or as a supplement is about 85% (Saubert et al 1987, Gregory 1989, 1995, 1997, Pfeiffer et al 1997, Cuskelly et al 1996). Folic acid as a supplement is almost 100% bioavailable on an empty stomach. Picciano et al (2004) have recently demonstrated that the inclusion of cows' milk in the diet enhances the bioavailability of food folate as assessed by changes in erythrocyte folate and plasma total homocysteine concentrations, but not when assessed by plasma folate concentrations. Some controlled studies to assess requirements have used a defined diet containing food folate and supplemented with folic acid, so the term dietary folate equivalents (DFE) has been used to accommodate the varying bioavailabilities.

1 µg dietary folate equivalent (DFE)	= 1 µg food folate
	= 0.5 µg folic acid on an empty stomach
	= 0.6 µg folic acid with meals or as fortified foods

Inadequate folate intake leads to decreased serum folate, then decreased erythrocyte folate, a rise in homocysteine and megaloblastic changes in bone marrow and other rapidly dividing tissues (Eichner & Hillman 1971). As depletion progresses, macrocytic cells are produced and macrocytic anaemia develops. Eventually, full-blown anaemia results in weakness, fatigue, irritability and palpitations. Folic acid supplementation in pregnancy can reduce both the occurrence and recurrence of neural tube defects in the newborn (Bower & Stanley 1989, CDC 1992, Czeizel & Dudas 1992, Kirke et al 1993, Laurence et al 1981, Wald et al 1991).

Indicators of folate requirement include erythrocyte, serum or urinary folate, plasma homocysteine and haematological status measures as well as clinical endpoints such as neural tube defects or chronic degenerative disease. Of these, erythrocyte folate is generally regarded as the primary indicator as it reflects tissue folate stores. For some age groups, erythrocyte folate is used in conjunction with plasma homocysteine and plasma or serum folate.

RECOMMENDATIONS BY LIFE STAGE AND GENDER

<i>Infants</i>	AI	Folate (as dietary folate equivalents)
0–6 months	65 µg/day (as folate)	
7–12 months	80 µg/day	

Rationale: The AI for 0–6 months was calculated by multiplying together the average intake of breast milk (0.78 L/day) and the average concentration of folate in breast milk of 85 µg/L (Asfour et al 1977, Ek & Magnus 1982, FNB:IOM 1998, Salmenpera et al 1986, Smith et al 1983, 1985), and rounding.

The AI for 7–12 months was set by the reference body weight ratio, estimating up from young infants or down from adults. Both estimates gave an AI of 80 µg/L which is also consistent with data for older, fully breast-fed or fully formula-fed infants in the studies of Asfour et al (1977), Ek & Magnus (1982), Salmenpera et al (1986) and Smith et al (1983).

<i>Children & adolescents</i>	EAR	RDI	Folate (as dietary folate equivalents)
All			
1–3 yr	120 µg/day	150 µg/day	
4–8 yr	160 µg/day	200 µg/day	
Boys			
9–13 yr	250 µg/day	300 µg/day	
14–18 yr	330 µg/day	400 µg/day	
Girls			
9–13 yr	250 µg/day	300 µg/day	
14–18 yr	330 µg/day	400 µg/day	

Rationale: As there are no experimental data for children, the EARs were set by extrapolation from adult data using metabolic body weight ratios with an allowance for growth as per FNB:IOM (1998). In the absence of information on the standard deviation of the requirement, the RDI was set assuming a CV of 10% for the EAR.

Adults	EAR	RDI	Folate (as dietary folate equivalents)
Men			
19–30 yr	320 µg/day	400 µg/day	
31–50 yr	320 µg/day	400 µg/day	
51–70 yr	320 µg/day	400 µg/day	
>70 yr	320 µg/day	400 µg/day	
Women			
19–30 yr	320 µg/day	400 µg/day	
31–50 yr	320 µg/day	400 µg/day	
51–70 yr	320 µg/day	400 µg/day	
>70 yr	320 µg/day	400 µg/day	

Rationale: The EAR for younger adults was set by reference to metabolic balance studies, notably the long term maintenance study in women that found no difference in mean final erythrocyte folate at 400 µg/day compared to 200–300 µg/day but a higher number of subjects with low erythrocyte folate, lower mean plasma folate and increased homocysteine levels (O’Keefe et al 1995). Other studies taken into account as cited in FNB:IOM (1998) were Herbert (1962a,b), Jacob et al (1994), Krumdieck et al (1978), Milne et al (1983), Sauberlich et al (1987), Stites et al (1997), von der Porten (1992) and Zalusky & Herbert (1961). For adults over 51 years, the requirements were based on metabolic, observational and epidemiological studies (Bates et al 1980, Garry et al 1982, Jagerstad 1977, Jagerstad & Westesson 1979, Koehler et al 1996, Ortega et al 1993, Rosenburg 1992, Sayoun 1992, Sayoun et al 1988, Selhub et al 1993, Tucker et al 1996, 1984).

In the absence of information on the SD of the requirement, the RDI was set assuming a CV of 10% for the EAR.

Special note: Evidence about the levels of folic acid needed in women to prevent neural tube defects did not form the basis for the adult EARs and RDIs. Women capable of, or planning, pregnancies should consume additional folic acid as a supplement or in the form of fortified foods at a level of 400 µg/day folic acid for at least one month before and three months after conception, in addition to consuming food folate from a varied diet.

Pregnancy	EAR	RDI	Folate (as dietary folate equivalents)
14–18 yr	520 µg/day	600 µg/day	
19–30 yr	520 µg/day	600 µg/day	
31–50 yr	520 µg/day	600 µg/day	

Rationale: Folate requirements increase substantially in pregnancy. This recommendation does not include consideration of additional needs to prevent neural tube defects as the neural tube is formed before most women know they are pregnant. The data indicate that maximal protection against NTD is obtained when the mother is consuming very high levels (5,000 µg) of folic acid as supplements, in the

month preceding conception and in the first trimester (Wald et al 2001). Recommendations are based on evidence from controlled metabolic studies (Caudill et al 1997) and a series of population studies (Chanarin et al 1968, Colman et al 1975, Dawson 1966, Hansen & Rybo 1967, Lowenstein et al 1966, Qvist et al 1986, Willoughby 1967, Willoughby & Jewel 1966). The RDI was estimated assuming a CV of 10% for the EAR.

Lactation	EAR	RDI	Folate (as dietary folate equivalents)
14–18 yr	450 µg/day	500 µg/day	
19–30 yr	450 µg/day	500 µg/day	
31–50 yr	450 µg/day	500 µg/day	

Rationale: To estimate total folate requirement for lactation, the amount needed to provide sufficient breast milk folate (including a 50% bioavailability correction factor) was added to the EAR for adult women using the formula $0.78 \text{ L (volume)} \times 85 \text{ µg/L (concentration)} \times 2 \text{ (for bioavailability)} = 133 \text{ µg/day (+ 320 µg/day)}$. The RDI was estimated assuming a CV of 10% for the EAR.

UPPER LEVEL OF INTAKE - DIETARY FOLATE EQUIVALENTS

Infants

0–12 months **Not possible to establish for supplemental folic acid.**
Source of intake should be milk, formula and food only

ULs from fortified foods or supplements

Children and adolescents

1–3 yr 300 µg/day as folic acid
 4–8 yr 400 µg/day as folic acid
 9–13 yr 600 µg/day as folic acid
 14–18 yr 800 µg/day as folic acid

Adults 19+ yr

Men 1,000 µg/day as folic acid
 Women 1,000 µg/day as folic acid

Pregnancy

14–18 yr 800 µg/day as folic acid
 19–50 yr 1,000 µg/day as folic acid

Lactation

14–18 yr 800 µg/day as folic acid
 19–50 yr 1,000 µg/day as folic acid

Rationale: No adverse effects have been associated with consumption of the amounts of dietary folate equivalents normally found in foods or fortified foods (Butterworth & Tamura 1989). High supplemental intakes of folic acid have been shown to be related to adverse neurological effects in people with B₁₂ deficiency as they can precipitate or exacerbate the B₁₂ deficiency (Israels & Wilkinson 1949, Schwartz et al 1950, Spies et al 1948, Will et al 1959). General toxicity (Hunter et al 1970), increased carcinogenesis (Selby et al 1989) and adverse reproductive and developmental effects have also been reported (Czeizel & Dudas 1992, Czeizel et al 1994, Holmes-Siedle et al 1992, Kirke et al 1992, Lawrence et al 1981, Mukerjee et al 1984, Smithells et al 1981, Vergel et al 1990, Wald et al 1991).

In line with the FNB:IOM (1998) findings, setting of the LOAEL was based on the neurological effects seen with B₁₂ deficiency, as this is a fairly common deficiency in the population and as these data have some dose-response characteristics. A LOAEL of 5 mg/day was set on the basis of the studies described above, as there were 100 cases of neurological damage above this level but only 8 below. A UF of 5 was used as the dose-response data were not well controlled, the adverse effects are severe and a LOAEL only, rather than a NOAEL, was available.

The UL was therefore estimated to be 1 mg folic acid (1,000 µg)/day for adults. There are no data to suggest increased susceptibility in pregnancy or lactation, so the adult UL was applied to these groups as well. There is little direct evidence for other ages, so the UL was set on a relative body weight basis for children and adolescents. It was not possible to set a UL for infants.

REFERENCES

- Asfour R, Wahbeh N, Waslien CI, Guindi S, Darby WJ. Folic acid requirement of children. III. Normal infants. *Am J Clin Nutr* 1977;30:1098–105.
- Australian Bureau of Statistics: Commonwealth Department of Health and Aged Care. *National Nutrition Survey. Nutrient intakes and physical measurements*. Australia, 1995. Canberra: Australian Bureau of Statistics, 1998.
- Bates CJ, Fleming M, Paul AA, Black AE, Mandal AR. Folate status and its relation to vitamin C in healthy elderly men and women. *Age Ageing* 1980;9:241–8.
- Bower C, Stanley FJ. Dietary folate as a risk factor for neural tube defects: evidence from a case-control study in Western Australia. *Med J Aust* 1989;150:613–9.
- Butterworth CE, Tamura T. Folic acid safety and toxicity: a brief review. *Am J Clin Nutr* 1989;50:353–8.
- Caudill MA, Cruz AC, Gregory JF 3rd, Hutson AD, Bailey LB. Folate status response to controlled folate intake in pregnant human subjects. *J Nutr* 1997;127:2363–70.
- CDC (Centres for Disease Control and Prevention). Recommendations for the use of folic acid to reduce the number of cases of spina bifida and other neural tube defects. *Morb Mortal Wkly Rep* 1992;41:1–7.
- Chanarin I, Rothman D, Ward A, Perry J. Folate status and requirement in pregnancy. *Br Med J* 1968;2:390–4.
- Colman N, Larsen JV, Barker M, Barker EA, Green M, Metz J. Prevention of folate deficiency by food fortification.3. Effect in pregnant subjects of varying amounts of added folic acid. *Am J Clin Nutr* 1975;28:465–70.
- Cuskelly GJ, McNulty HM, Scott JM. Effect of increasing dietary folate on red-cell folate: implications for prevention of neural tube defects. *Lancet* 1996;347:657–9.
- Czeizel AE, Dudas I. Prevention of the first occurrence of neural-tube defects by periconceptual vitamin supplementation. *N Engl J Med* 1992;327:1832–5.
- Czeizel AE, Dudas I, Metneki J. Pregnancy outcomes in a randomised controlled trial of periconceptual multivitamin supplementation. Final report. *Arch Gynecol Obstet* 1994;255:131–9.
- Dawson DW. Microdoses of folic acid in pregnancy. *J Obstet Gynaecol Br Commonw* 1966;73:44–8.
- Eichner ER, Hillman RS. The evolution of anemia in alcoholic patients. *Am J Med* 1971;50:218–32.
- Ek J, Magnus E. Plasma and red cell folate values and folate requirements in formula-fed term infants. *J Pediatr* 1982;100:738–44.
- Food and Nutrition Board: Institute of Medicine. *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline*. Washington DC: National Academy Press, 1998.

- Garry PJ, Goodwin JS, Hunt WC, Hooper EM, Leonard AG. Nutritional status in a healthy elderly population: dietary and supplemental intakes. *Am J Clin Nutr* 1982;36:319–31.
- Garry PJ, Goodwin JS, Hunt WC. Folate and vitamin B₁₂ status in a healthy elderly population. *J Am Geriatr Soc* 1984;32:719–26.
- Gregory JF 3rd. The bioavailability of folate. In: Bailey LB, ed. *Folate in health and disease*. New York: Marcel Dekker, 1995. Pp195–235.
- Gregory JF 3rd. Bioavailability of folate. *Eur J Clin Nutr* 1997;51:S54–S59.
- Gregory JF 3rd. Chemical and nutritional aspects of folate research: analytical procedures, methods of folate synthesis, stability and bioavailability of dietary folates. *Adv Food Nutr Res* 1989;33:1–101.
- Hansen H, Rybo G. Folic acid dosage in prophylactic treatment during pregnancy. *Acta Obstet Gynecol Scand* 1967;46:107–12.
- Herbert V, Das KC. Folic acid and vitamin B₁₂. In: Shils ME, Olson JA, Shike M. eds. *Modern nutrition in health and disease, 8th edition*. Philadelphia: Lea & Febiger, 1993. Pp 402–25.
- Herbert V. Experimental nutritional folate deficiency in man. *Trans Assoc Am Physicians* 1962a;75:307–20.
- Herbert V. Minimal daily adult folate requirement *Arch Intern Med* 1962b;110:649–52.
- Holmes-Siedle M, Lindenbaum RH, Galliard A. Recurrence of neural tube defect in a group of at risk women: a ten year study of Pregnavite Forte F. *J Med Genet* 1992;29:134–5.
- Hunter R, Barnes J, Oakeley HF, Matthews DM. Toxicity of folic acid given in pharmacological doses to healthy volunteers. *Lancet* 1970;1:61–3.
- Israels MC, Wilkinson JF. Risk of neurological complications in pernicious anemia treated with folic acid. *Br Med J* 1949;2:1072–5.
- Jacob RA, Wu M-M, Henning SM, Swendseid ME. Homocysteine increases as folate decreases in plasm of healthy men during short term dietary folate and methyl group restriction. *J Nutr* 1994;124:1072–80.
- Jagerstad M, Westesson A-K. Folate. *Scand J Gastroentero* 1979;14:196–202.
- Jagerstad M. Folate intake and blood folate in elderly subjects, a study using the double sampling portion technique. *Nutr Metab* 1977;21:29–31.
- Kirke PN, Daly LE, Elwood JH. A randomised trial of low-dose folic acid to prevent neural tube defects. *Arch Dis Child* 1992;67:1442–6.
- Kirke PN, Molloy AM, Daly LE, Burke H, Weir DG, Scott JM. Maternal plasma folate and vitamin B₁₂ are independent risk factors for neural tube defects. *Q J Med* 1993;86:703–8.
- Koehler KM, Romero LJ, Stauber PM, Pareo-Tubbeh SL, Liang HC, Baumgartner RN, Garry PJ, Allen RH, Stabler SP. Vitamin supplementation and other variables affecting serum homocysteine and methylmalonic acid concentrations in elderly men and women. *J Am Coll Nutr* 1996;15:364–76.
- Krumdieck CL, Fukushima K, Fukushima T, Shiota T, Butterworth CE Jnr. A long-term study of the excretion of folate and pterins in a human subject after ingestion of ¹⁴C folic acid, with observations on the effect of diphenylhydantoin administration. *Am J Clin Nutr* 1978;31:88–93.
- Laurence KM, James N, Miller MH, Tennant GB, Campbell H. Double-blind randomised controlled trial of folate treatment before conception to prevent recurrence of neural tube defects. *Br Med J* 1981;282:1509–11.
- Lawrence VA, Lowenstein JE, Eichner ER. Aspirin and folate binding; in vivo and in vitro studies of serum binding and urinary excretion of endogenous folate. *J Lab Clin Invest* 1981;103:944–8.
- Lowenstein I, Cantlie G, Ramos O, Brunton L. The incidence and prevention of folate deficiency in pregnant clinic population. *Can Med Assoc J* 1966;95:797–806.

- Milne DB, Johnson K, Mahalko JR, Sandstead HH. Folate status of adult males living in a metabolic unit; possible relationships with iron nutriture. *Am J Clin Nutr* 1983;37:768–73.
- Ministry of Health. NZ food: NZ People. Key results of the 1997 National Nutrition Survey. Wellington: Ministry of Health, 1999.
- Mukerjee MD, Sandstead HH, Ratnaparkhi MV, Johnson LK, Milne DB, Stelling HP. Maternal zinc, iron, folic acid and protein nutriture and outcome of human pregnancy. *Am J Clin Nutr* 1984;40:496–507
- O'Keefe CA, Baily LB, Thomas EA, Hofler SA, Davis BA, Cerda JJ, Gregory JF 3rd. Controlled dietary folate affects folate status in nonpregnant women. *J Nutr* 1995;125:2717–25.
- Ortega RM, Redondo R, Andres P, Equleor I. Nutritional assessment of folate and cyanocobalamin status in a Spanish elderly group. *Int J Vitam Nutr Res* 1993;63:17–21.
- Pfeiffer CM, Rogers LM, Bailey LB, Gregory JF. Absorption of folate from fortified cereal-grain products and of supplemental folate consumed with or without food determined by using a dual-label stable-isotope protocol. *Am J Clin Nutr* 1997;66:1388–97.
- Picciano MF, West SG, Ruch AL, Kris-Etherton PM, Zhao G, Johnston KE, Maddox DH, Fishell VK, Dirienzo DB, Tamura T. Effect of cow milk on food folate bioavailability in young women. *Am J Clin Nutr* 2004;80:1565–9.
- Qvist I, Abdulla M, Jagerstad M, Svensson S. Iron, zinc and folate status during pregnancy and two months after delivery. *Acta Obstet Gynecol Scand* 1986;65:15–22.
- Rosenburg IH. Folate. In: Hartz SDC, Russell RM, Rosenberg IH, eds. *Nutrition in the elderly. The Boston Nutritional Status Survey*. London: Smith-Gordon, 1992. Pp 135–9.
- Salmenpera L, Perheentupa J, Simes MA. Folate nutrition is optimal in exclusively breast-fed infants but inadequate in some of the mothers and in formula-fed infants. *J Pediatr Gastroenterol Nutr* 1986;5:283–9.
- Sauberlich HE, Kretsch MJ, Skala JH, Johnson HL, Taylor PC. Folate requirement and metabolism in nonpregnant women. *Am J Clin Nutr* 1987;46:1016–28.
- Sayoun NR, Otradovec CL, Hartz SC, Jacob RA, Peters H, Russell RM, McGandy RB. Dietary intakes and biochemical indicators of nutritional status in an elderly institutionalised population. *Am J Clin Nutr* 1988;47:524–33.
- Sayoun N. Nutrient intake by the NNS elderly population. In: Hartz SDC, Russell RM, Rosenberg IH, eds. *Nutrition in the elderly. The Boston Nutritional Status Survey*. London: Smith-Gordon, 1992. Pp 31–44.
- Schwartz SO, Kaplan SR, Armstrong BE. The long-term evaluation of folic acid in the treatment of pernicious anemia. *J Lab Clin Med* 1950;35:894–8.
- Selby JV, Friedman GD, Fireman BH. Screening prescription drugs for possible carcinogenicity: eleven to fifteen years of follow-up. *Cancer Res* 1989;49:5736–47.
- Selhub J, Jacques PF, Wilson PWF, Rush D, Rosenberg IH. Vitamin status and intake as primary determinants of homocysteinemia in an elderly population. *JAMA* 1993;270:2693–8.
- Smith AM, Picciano MF, Deering RH. Folate supplementation during lactation: maternal folate status, human milk folate content and their relationship to infant folate status. *J Pediatr Gastroenterol Nutr* 1983;2:622–8.
- Smith AM, Picciano MF, Deering RH. Folate intake and blood concentrations of term infants. *Am J Clin Nutr* 1985;41:590–8.
- Smithells RW, Sheppard S, Schorah CJ, Sellar MJ, Nevin NC, Harris R, Reads AP, Fielding DW. Apparent prevention of neural tube defects by periconceptual vitamin supplementation. *Arch Dis Child* 1981;56:911–8.

- Spies TD, Stone RE, Lopez GG, Milanese F, Aramburu T, Toca RL. The association between gastric achlorhydria and subacute degeneration of the spinal cord. *Postgrad Med* 1948;4:89–95.
- Stites TE, Bailey LB, Scott KC, Toth JP, Fisher WP, Gregory JF 3rd. Kinetic modelling of folate metabolism through use of chronic administration of deuterium-labelled folic acid in men. *Am J Clin Nutr* 1997;65:53–60.
- Tucker KL, Selhub J, Wilson PZW, Rosenberg H. dietary intake pattern relates to plasma folate and homocysteine concentrations in the Framingham Heart Study. *J Nutr* 1996;126:3025–31.
- van der Put NM, Steegers-Theunissen RP, Frosst P, Trijbels FJ, Eskes TK, van den Heuvel LP, Mariman EC, den Heijer M, Rozen R, Blom HJ. Mutated methylene-tetrahydrofolate reductase as a risk factor for spina bifida. *Lancet* 1995;346:1070–1.
- Vergel RG, Sanchez LR, Heredero BL, Rodriguez PL, Martinez AJ. Primary prevention of neural tube defects with folic acid supplementation: Cuban experience. *Prenat Diagn* 1990;10:149–52.
- Von der Porten AE, Gregory JF 3rd, Toth JP, Cerda JJ, Curry SH, Bailey LB. In vivo folate kinetics during chronic supplementation of human subjects with deuterium-labelled folic acid. *J Nutr* 1992;122:1293–9.
- Wagner C. Symposium on the subcellular compartmentalisation of folate metabolism. *J Nutr* 1996;126:1228S–1234S.
- Wald N, Sneddon J, Densem J, Frost C, Stone R. Prevention of neural tube defects: results of the Medical Research Council vitamin study. *Lancet* 1991;338:131–7.
- Wald NJ, Law MR, Morris JK, Wald DS. Quantifying the effect of folic acid. *Lancet* 2001;358:2069–73.
- Whitehead VM. Pharmacokinetics and physiological disposition of folate and its derivatives. In: Blakely RL, Whitehead VM, eds. *Folates and pterins, Vol 3*. New York: John Wiley & Sons, 1986. Pp 177–205.
- Will JJ, Mueller JF, Brodine C, Kiely CE, Friedman B, Hawkins VR, Duta J, Vilter RN. Folic acid and vitamin B12 in pernicious anemia. Studies on patients treated with these substances over a ten-year period. *J Lab Clin Med* 1959;53:22–38.
- Willoughby ML, Jewel FJ. Investigation of folic acid requirements in pregnancy. *Br Med J* 1966;2:1568–71.
- Willoughby ML. An investigation of folic acid requirements in pregnancy. II. *Br J Haematol* 1967;13:503–9.
- Zalusky R, Herbert V. Megaloblastic anemia in scurvy with response to 50 micrograms of folic acid daily. *N Eng J Med* 1961;265:1033–8.