

The following is an extract from:

Nutrient Reference Values for Australia and New Zealand
Including Recommended Dietary Intakes

ENDORSED BY THE NHMRC ON 9 SEPTEMBER 2005

© Commonwealth of Australia 2006

ISBN Print 1864962372
ISBN Online 1864962437

The Nutrient Reference Values (NRVs) was a joint initiative of the Australian National Health and Medical Research Council (NHMRC) and the New Zealand Ministry of Health (MoH). The NHMRC would like to thank the New Zealand MoH for allowing the use of the NRV material in the development of this website.

NHMRC publications contact:

Email: nhmrc.publications@nhmrc.gov.au
Internet: <http://www.nhmrc.gov.au>
Free Call: 1800 020 103 ext 9520

VITAMIN E

BACKGROUND

Vitamin E is the name given to a group of water-insoluble, plant-derived substances. There are eight naturally-occurring isomers and a number of semisynthetic or synthetic homologues. The naturally-occurring *d*- (or RRR) alpha-tocopherol is the most biologically active form and vitamin E activity is traditionally expressed in terms of equivalents of this isomer (mg alpha-tocopherol equivalents or α -TE). Other tocopherols such as gamma-tocopherol also have vitamin E activity. There are four tocopherol homologues (*d*- α -, *d*- β -, *d*- γ - and *d*- δ -) and four tocotrienols. Other forms of vitamin E occur in lower amounts in foods and are less active in animal bioassay. The usual form in supplements is synthetic *dl*- (or all-*rac*) α -tocopherol that consists of a mixture of active and inactive stereoisomers, because natural vitamin E from wheat germ oil is expensive. The equivalence of the various forms is shown below:

Form	Alternative name	mg α -tocopherol equivalence
<i>d</i> - α -tocopherol	RRR- α -tocopherol	1
<i>d</i> - α -tocopherol acetate	RRR- α -tocopherol acetate	0.91
<i>d</i> - α -tocopherol acid succinate	RRR- α -tocopherol acid succinate	0.81
<i>dl</i> - α -tocopherol	all- <i>rac</i> - α -tocopherol	0.74
<i>dl</i> - α -tocopherol acetate	all- <i>rac</i> - α -tocopherol	0.67
<i>d</i> - β -tocopherol	RRR- β -tocopherol	0.25–0.40
<i>d</i> - γ -tocopherol	RRR- γ -tocopherol	0.10
α -tocotrienol		0.25–0.30

The major role of vitamin E is to protect polyunsaturated fatty acids (PUFA) from oxidation. It acts as an anti-oxidant in the lipid phase of cell membranes. Vitamin E requirements have been reported to increase when intakes of PUFA are increased (Dam 1962, Horwitt 1962) and a ratio of at least 0.4 mg α -tocopherol per gram of PUFA has been recommended (Bieri & Evarts 1973, Horwitt 1974, Witting & Lee 1975). Most dietary sources of polyunsaturated fat are also relatively rich in vitamin E. However supplements of fish oils or other pure n-3 fatty acids may not provide the amount of vitamin E needed.

The activity vitamin E complements that of selenium-dependent glutathione peroxidase in protecting the membrane PUFAs from free radical damage. Although vitamin E is mainly located in cells and organelle membranes, its concentration may be very low, suggesting that after its reaction with free radicals it is rapidly regenerated, possibly by other antioxidants such as selenium, ubiquinol and vitamin C (Doba et al 1985, Niki et al 1982, Stoyanovsky et al 1995).

The main source of vitamin E is fats and oils. It is also found in some vegetables, in the fats of meat, poultry and fish and, to lesser degrees, in cereals and dairy foods. About half the tocopherol in wheat germ, sunflower, safflower, canola, olive and cottonseed oils is α -tocopherol but soybean and corn oils contain about 10 times as much γ -tocopherol as α -tocopherol. Most vitamin E is found in foods containing fat. Absorption requires micelle formation and chylomicron secretion in the gut (Muller et al 1974) together with biliary and pancreatic secretions. Efficiency of absorption is low, but the precise rate is unknown.

Vitamin E is transported in the blood by the plasma lipoproteins and erythrocytes. Tocopherols are carried from the gut to the liver in chylomicrons where they are incorporated as chylomicron remnants. Catabolism of chylomicrons takes place in the systemic circulation through the action of cellular lipoprotein lipase. Vitamin E can be transferred to high density lipoprotein (HDL) and then

to low density lipoprotein (LDL) and very low density lipoprotein (VLDL). Most α -tocopherol enters peripheral tissues within the intact lipoprotein through the LDL receptor pathway.

Although all tocopherol homologues are absorbed similarly, α -tocopherol predominates in blood and tissue as the binding proteins take it up preferentially. Plasma vitamin E and tissue concentrations vary little over a wide range of dietary intake and the brain is particularly resistant to depletion (Bourne & Clement 1991).

The main oxidation product of α -tocopherol is tocopheryl quinone which is conjugated to glucuronate and is excreted in bile or further degraded in the kidneys to α -tocopheronic acid before excretion in bile (Drevon 1991). Some may be excreted through the skin (Shiratori 1974).

Overt deficiency symptoms in normal individuals consuming diets low in vitamin E have never been described. It occurs only as a result of genetic abnormalities, fat malabsorption syndrome (Rader & Brewer 1993, Sokol 1993) or protein-energy malnutrition (Kalra et al 1998, Laditan & Ete 1982). The main symptom is a peripheral neuropathy. Other symptoms include spinocerebellar ataxia, skeletal myopathy and pigmented retinopathy (Sokol 1988).

In epidemiological studies, higher intakes of vitamin E have been related to reduction in cardiovascular disease risk (Gey et al 1991, Rimm & Stampfer 1993, Stampfer et al 1993), diabetic complications (Baynes 1991, Mullarkey et al 1990, Semenkovich & Heinecke 1997), certain cancers (Comstock et al 1997, Eichhlozer et al 1996, Yong et al 1997) and cataracts (Jacques & Chylack 1991, Knekt et al 1992, Leske et al 1991). Not all studies, however, have confirmed a relationship and clinical trials with supplements in high risk groups, have shown little benefit. Further discussion of these trials is given in the 'Chronic disease' section.

Indicators that have been used to estimate vitamin E requirements include lipid peroxidation markers, oxidation products of DNA or proteins, vitamin E metabolite excretion, vitamin E biokinetics, vitamin E deficiency symptoms, plasma α -tocopherol concentration, hydrogen peroxide-induced haemolysis or the relationship of vitamin E to chronic disease status. However, erythrocyte fragility studies have been the most widely used.

The US DRI review in 2000 used the data of Horwitt (1960, 1963). These same data had been used in setting the earlier US RDIs but were interpreted differently in 2000, leading to considerably increased recommendations. In the US DRI review of 2000, the amount of dietary vitamin E required to bring plasma α -tocopherol to a level where per cent haemolysis was low was used to estimate an EAR (Horwitt 1960, 1963). However, the interpretation of these data is problematic in relation to level of plasma α -tocopherol at which adverse effects are seen, as there were no data available for plasma α -tocopherol concentrations between 5 and 12 $\mu\text{mol/L}$. All four subjects below 6 $\mu\text{mol/L}$ plasma α -tocopherol (range 2–5 $\mu\text{mol/L}$) had haemolysis of about 80% or above and all 6 subjects with concentrations between 12 and 22 $\mu\text{mol/L}$, had haemolysis of 12% or less. There has been disagreement as to whether the 'adequacy' cut off should be midway between these two clusters or at the lowest point showing low haemolysis. The data are dichotomous, not continuous, thus preventing an accurate dose-response analysis. Changing the cut-off point makes a large difference to the estimated requirement. In addition, the authors of the key paper themselves expressed concern about the validity of the technique for assessing vitamin E requirements (Horwitt 1960, 1963, 2001).

Given these uncertainties, an AI rather than an EAR was set for vitamin E based on median population intakes in Australia and New Zealand – both healthy populations with no apparent vitamin E deficiency. Recommendations for infants were based on the median concentration in breast milk of healthy mothers.

RECOMMENDATIONS BY LIFE STAGE AND GENDER

<i>Infants</i>	AI	Vitamin E (as α-tocopherol equivalents)
0–6 months	4 mg/day	
7–12 months	5 mg/day	

Rationale: The AI for 0–6 months was calculated by multiplying the average intake of breast milk (0.78 L/day) by the average concentration of vitamin E in breast milk of 4.9 mg/L (Boersma et al 1991, Chappell et al 1985, Jansson et al 1981, Lammi-Keefe et al 1985, 1990) and rounding. Two of these studies reported only α -tocopherol data but Boersma et al (1991) showed that the tocopherol content of breast milk is almost entirely comprised of α -tocopherol. For 7–12 months, the AI was extrapolated from younger infants on a body weight basis and rounded.

<i>Children & adolescents</i>	AI	Vitamin E (as α-tocopherol equivalents)
All		
1–3 yr	5 mg/day	
4–8 yr	6 mg/day	
Boys		
9–13 yr	9 mg/day	
14–18 yr	10 mg/day	
Girls		
9–13 yr	8 mg/day	
14–18 yr	8 mg/day	

Rationale: As there are no specific data on which to base an EAR for children and adolescents, an AI was set based on the median intakes in Australia and New Zealand from the National Nutrition Surveys with rounding up to the nearest milligram (ABS 1998, MOH 1999, 2003).

<i>Adults</i>	AI	Vitamin E (as α-tocopherol equivalents)
Men		
19–30 yr	10 mg/day	
31–50 yr	10 mg/day	
51–70 yr	10 mg/day	
>70 yr	10 mg/day	
Women		
19–30 yr	7 mg/day	
31–50 yr	7 mg/day	
51–70 yr	7 mg/day	
>70 yr	7 mg/day	

Rationale: As there are not sufficient data on which to base an EAR for adults, an AI was set based on the median intakes in Australia and New Zealand from the National Nutrition Surveys with rounding up to the nearest milligram (ABS 1998, MOH 1999). The values set for men and women were the highest median intake for any respective adult age band.

Pregnancy	AI	Vitamin E (as α-tocopherol equivalents)
14–18 yr	8 mg/day	
19–30 yr	7 mg/day	
31–50 yr	7 mg/day	

Rationale: There is no evidence of increased needs for vitamin E in pregnancy, so the AI is set at that for the non-pregnant woman.

Lactation	AI	Vitamin E (as α-tocopherol equivalents)
14–18 yr	12 mg/day	
19–30 yr	11 mg/day	
31–50 yr	11 mg/day	

Rationale: The AI for lactation is set at that for the non-lactating woman plus an allowance for the vitamin E secreted in milk.

UPPER LEVEL OF INTAKE - VITAMIN E - (as α -tocopherol equivalents)

Infants

0–12 months **Not possible to establish. Source of intake should be breast milk, formula and food only**

Children

1–3 yr 70 mg/day
4–8 yr 100 mg/day

Boys

9–13 yr 180 mg/day
14–18 yr 250 mg/day

Girls

9–13 yr 180 mg/day
14–18 yr 250 mg/day

Adults 19+ yr

Men 300mg/day
Women 300mg/day

Pregnancy

All ages 300 mg/day

Lactation

All ages 300 mg/day

Rationale: In recent years, several clinical intervention trials have assessed the effects of high doses of vitamin E on chronic disease outcomes, including the CHAOS Heart trial which used 268–567 mg *d*- α -tocopherol/day (Stephens et al 1996), the GISSI study with 300 mg vitamin E as synthetic α -tocopherol (GISSI-Prevenzione Investigators 1999), the ATBC study using 55 mg *dl* α -tocopherol (ATBC 1994, Heinonen et al 1998), the HOPE study using 268 mg vitamin E (Yusuf et al 2000), the Primary Prevention Study using 300 mg/day synthetic α -tocopherol (Collaborative group of the Primary Prevention Study 2001) and the Heart Protection Study with 600 mg of vitamin E (Heart Protection Study Collaborative Group, 2002). In addition, there have been a number of experimental trials using supplements ranging from 540 to 970 mg *d*- α -TEs. With the exception of an increase in subarachnoid haemorrhaging in smoking hypertensives in the ATBC study (Leppanen et al 2000a,b), a non-significant increase in stroke (relative risk 1.17) in the HOPE study and a tendency to haemorrhage in aspirin users in the Primary Prevention Project, no adverse events have been recorded. However, most studies were not specifically designed to assess adverse events to Vitamin E alone.

Meydani et al (1998) undertook an experimental, dose-dependent study in 88 healthy volunteers aged >65 years, with one control group and three varying dose groups (equivalent to 34, 134 or 537 mg *d*- α -TEs), over 4 months. This study had the most comprehensive assessment of potential adverse events. There were no subjective side effects and no effects on glutathione peroxidase, superoxide dismutase, immunoglobulin, anti-DNA or anti-thyroglobulin antibodies, body weight, total plasma proteins, albumin, glucose, lipids or lipoprotein profile, total bilirubin, serum liver enzymes, blood count, platelet number, bleeding time, haemoglobin, haematocrit or urinary or serum creatinine. The NOAEL established from this study was 540 mg/day. A UF of 2 was applied to cover inter-individual differences in sensitivity. A larger UF was not considered necessary because data from a number of other less well controlled studies showed no adverse effects at considerably higher intakes. The UL for vitamin E was therefore established as 270 mg/day for adults and rounded to 300 mg/day. The ULs for other age groups were derived on a relative body weight basis.

REFERENCES

- ATBC The Alpha-Tocopherol, Beta-Carotene Cancer Prevention Study Group. The effect of vitamin E and beta-carotene on the incidence of lung cancer and other cancers in male smokers. *N Engl J Med* 1994;330:1029–35.
- Australian Bureau of Statistics/Commonwealth Department of Health and Ageing. *National Nutrition Survey: Nutrient intakes and physical measurements. Australia, 1995*. Canberra: Australian Bureau of Statistics, 1998.
- Baynes JW. Role of oxidative stress in development of complications in diabetes. *Diabetes* 1991;40:405–12.
- Bieri JG, Everts RP. Gamma-tocopherol: metabolism, biological activity and significance in human vitamin E nutrition. *Am J Clin Nutr* 1973;27:980–6.
- Boersma ER, Offringa PJ, Muskiet FA, Chase WM, Simmons IJ. Vitamin E, lipid fractions and fatty acid composition of colostrum, transitional milk and mature milk: an international comparative study. *Am J Clin Nutr* 1991;53:1197–204.
- Bourne J, Clement M. Kinetics of rat peripheral nerve, forebrain and cerebellum α -tocopherol depletion: comparison with different organs. *J Nutr* 1991;121:1204–7.
- Chappell JE, Francis T, Clandinin MT. Vitamin A and E content of human milk at early stages of lactation. *Early Hum Devel* 1985;11:157–67.
- Collaborative Group of the Primary Prevention Project (PPP). Low-dose aspirin and vitamin E in people at cardiovascular risk: a randomized trial in general practice. *Lancet* 2001;357:89–95.

- Comstock GW, Alberg AJ, Huang HY, Wu K, Burke AE, Hoffman SC, Norkus EP, Gross M, Cutler RG, Morris JS, Spate VL, Helzlsouer KJ. The risk of developing lung cancer associated with antioxidants in the blood; ascorbic acid, carotenoids, alpha-tocopherol, selenium and total peroxy radical absorbing capacity. *Cancer Epidemiol Biomarkers Prev* 1997;6:907–16.
- Dam, H. Interrelations between vitamin E and polyunsaturated fatty acids in animals. *Vitam Horm* 1962;20:527–40.
- Doba T, Burton GW, Ingold KU. Antioxidant and co-antioxidant activity of vitamin C. The effect of vitamin C either alone or in the presence of vitamin E or a water-soluble vitamin E analogue, upon the peroxidation of aqueous multilamellar phospholipid liposomes. *Biochim Biophys Acta* 1985;835:298–303.
- Drevon CA. Absorption, transport and metabolism of vitamin E. *Free Radic Res Commun* 1991;14:229–46.
- Eichholzer M, Stahelin HB, Gey KF, Ludin E, Bernasconi F. Prediction of male cancer mortality by plasma levels of interacting vitamins: 17-year follow-up of the prospective Basel study. *Int J Cancer* 1996;66:1490S–1500S.
- Food and Nutrition Board: Institute of Medicine. *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium and Carotenoids*. Washington, DC: National Academy Press, 2000.
- Gey KF, Puska P, Jordan P, Moser UK. Inverse correlation between plasma vitamin E and mortality from ischaemic heart disease in cross-cultural epidemiology. *Am J Clin Nutr* 1991;53:326S–334S.
- GISSI-Prevenzione Investigators. Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. *Lancet* 1999;354:447–55.
- Heart Protection Study Collaborative Group. MRC/BHF Heart protection study of antioxidant vitamin supplementation in 20,536 high-risk individuals: a randomized, placebo-controlled trial. *Lancet* 2002;360:23–33.
- Heinonen OP, Albanes D, Virtamo J, Taylor PR, Huttunen JK, Hartman AM, Haapakoski J, Malila N, Rautalahti M, Ripatti S, Maenpaa H, Teerenhovi L, Koss L, Virolainen M, Edwards BK. Prostate cancer and supplementation with alpha-tocopherol and beta-carotene: incidence and mortality in a controlled trial. *J Natl Cancer Inst* 1998;90:440–6.
- Horwitt MK. Vitamin E and lipid metabolism in man. *Am J Clin Nutr* 1960;8:451–61.
- Horwitt, MK. Interrelations between vitamin E and polyunsaturated fatty acids in adult men. *Vitam Horm* 1962;20:541–58.
- Horwitt, MK. Status of human requirements for vitamin E. *Am J Clin Nutr* 1974;27:1182–93.
- Horwitt MK. Critique of the requirement for vitamin E. *Am J Clin Nutr* 2001;73:1003–5.
- Horwitt MK, Century B, Zeman AA. Erythrocyte survival time and reticulocyte levels after tocopherol depletion in man. *Am J Clin Nutr* 1963;12:99–106.
- Jacques PF, Chylack LT Jr. Epidemiologic evidence of a role for the antioxidant vitamins and carotenoids in cataract prevention. *Am J Clin Nutr* 1991;53:352S–355S.
- Jansson L, Akesson B, Holmberg L. Vitamin E and fatty acid composition of human milk. *Am J Clin Nutr* 1981;34:8–13.
- Kalra V, Grover J, Ahuja GK, Rathi S, Khurana DS. Vitamin E deficiency and associated neurological deficits in children with protein-energy malnutrition. *J Trop Pediatr* 1998;44:291–5.
- Knekt P, Heliovaara M, Rissanen A, Aromaa A, Aaran RK. Serum antioxidant vitamins and risk of cataract. *Br Med J* 1992;305:1392–4.
- Laditan AA, Ette SI. Plasma alpha-tocopherol (vitamin E) levels and tocopherol-lipid ratio among children with protein-energy malnutrition (PEM). *Ann Trop Paediatr* 1982;2:85–8.

- Lammi-Keefe CJ, Jensen RJ, Clark RM, Ferris AM. Alpha tocopherol, total lipid and linoleic acid contents of human milk at 2, 6, 12 and 16 weeks. In: Schaub J, ed. *Composition and physiological properties of human milk*. New York: Elsevier Science, 1985. Pp 241–5.
- Lammi-Keefe CJ, Ferris AM, Jensen RG. Changes in human milk at 0600, 1000, 1400, 1800 and 2200 h. *J Pediatr Gastroenterol Nutr* 1990;11:83–8.
- Leppala JM, Virtamo J, Fogelholm R, Albanes D, Taylor PR and Heinonen OP. Vitamin E and beta-carotene supplementation in high risk for stroke: a sub-group analysis of the Alpha-tocopherol, Beta-carotene Cancer Prevention Study. *Arch Neurol* 2000a;57:1503–9.
- Leppala JM, Virtamo J, Fogelholm R, Huttenen JK, Albanes D, Taylor PR, Heinonen OP. Controlled trial of alpha-tocopherol and beta-carotene supplements on stroke incidence and mortality in male smokers. *Arterioscler Thromb Vasc Biol* 2000b;20:230–5.
- Leske MC, Chylack LT Jr, Wu SY. The lens opacities case-control study. Risk factors for cataract. *Arch Ophthalmol* 1991;109:244–51.
- Meydani SN, Meydani M, Blumberg JB, Leka LS, Pedrosa M, Diamond R, Schaefer EJ. Assessment of the safety of supplementation with different amounts of vitamin E in healthy older adults. *Am J Clin Nutr* 1998;68:311–8.
- Ministry of Health. *NZ Food: NZ People. Key results of the 1997 National Nutrition Survey*. Wellington: Ministry of Health, 1999.
- Ministry of Health. *NZ Food: NZ Children. Key results of the 2002 National Children's Nutrition Survey*. Wellington: Ministry of Health, 2003.
- Mullarkey CJ, Edelstein D, Brownlee M. Free radical generation by early glycation products: a mechanism for accelerated atherogenesis in diabetes. *Biochem Biophys Res Commun* 1990;173:932–9.
- Muller DP, Harries JT, Lloyd JK. The relative importance of the factors involved in the absorption of vitamin E in children. *Gut* 1974;15:966–71.
- Niki E, Tsuchiya J, Tanimura R, Kamiya Y. Regeneration of vitamin E from alpha-chromanoxyl radical by glutathione and vitamin C. *Chem Lett* 1982;6:789–92.
- Rader JD, Brewer HB. Abetalipoproteinemia. New insights into lipoprotein assembly and vitamin E metabolism from a rare genetic disease *JAMA* 1993;270:865–9.
- Rimm EB, Stampfer MJ, Ascherio A, Giovannucci E, Colditz GA, Willett WC. Vitamin E consumption and the risk of coronary heart disease in men. *N Engl J Med* 1993;328:1450–6.
- Semenkovich CF, Heinecke JW. The mystery of diabetes and atherosclerosis: time for a new plot. *Diabetes* 1997;46:327–34.
- Shiratori T. Uptake, storage and excretion of chylomicron-bound 3H-alpha-tocopherol by the skin of the rat. *Life Sci* 1974;14:929–35.
- Sokol RJ. Vitamin E deficiency and neurologic disease. *Ann Rev Nutr* 1988;8:351–73.
- Sokol RJ. Vitamin E deficiency and neurological disorders. In : Packer I, Fuchs J, eds. *Vitamin E in health and disease*. New York: Marcel Dekker, 1993. Pp 815–49.
- Stampfer MJ, Hennekens CH, Manson JE, Colditz GA, Rosner B, Willett WC. Vitamin E consumption and the risk of coronary disease in women. *N Engl J Med* 1993;328:1444–9.
- Stephens NG, Parsons A, Schofield PM, Kelly F, Cheeseman K, Mitchinson MJ. Randomised controlled trial of vitamin E in patients with coronary disease: Cambridge Heart Antioxidant Study. *Lancet* 1996;347:781–5.
- Stoyanovsky DA, Osipov AN, Quinn PJ, Kagan VE. Ubiquinone-dependent recycling of vitamin E radicals by superoxide. *Arch Biochem Biophys* 1995;323:343–51.

- Witting LA, Lee L. Dietary levels of vitamin E and polyunsaturated fatty acids and plasma vitamin E. *Am J Clin Nutr* 1975;28:571–6.
- Yong LC, Brown CC, Schatzkin A, Dresser CM, Sleinski MJ, Cox CS, Taylor PR. Intake of vitamins E, C, and A and risk of lung cancer. The NHANES I epidemiologic follow-up study. *Am J Epidemiol* 1997;146:231–43.
- Yusuf S, Dagenais G, Pogue J, Bosch J, Sleight, P. Vitamin E supplementation and cardiovascular events in high-risk patients. The Heart Outcomes Prevention Evaluation Study Investigators. *New England Journal of Medicine* 2000;342:154–60.